

yellow. The venter is yellowish and its eyes are a silvery colour (Fig. 1). The snake is currently maintained as a captive in the Herpetology Laboratory at Instituto Butantan, São Paulo, Brazil.

REFERENCES

- Amaral, A. (1927a). Da ocorrência de albinismo em Cascavel, *Crotalus terrificus*. *Rev. Mus. Paul.* **15**, 55–61.
- Amaral, A. (1927b). Albinismo em “Dorme-dorme” (*Sibynomorphus turgidus*). *Rev. Mus. Paul.* **15**, 61–63.
- Amaral, A. (1932). Notas sobre chromatismo de ophidios – II. Casos de variação de colorido de certas serpentes. *Mem. Inst. Butantan* **Tomo VII**, 81–87.
- Amaral, A. (1934). Notas sobre chromatismo de ophidios – III. Um caso de xanthismo e um novo caso de albinismo, observados no Brasil. *Mem. Inst. Butantan*, **Tomo VIII**, 151–153.
- Andrade, D.V. & Abe, A.S. (1998). Abnormalities in a litter of *Boa constrictor amarali*. *The Snake* **28**, 28–32.
- Duarte, M.R.D.; Franco, F.L. & Oliveira, A.L. (2005). New records of albinism in neotropical snakes. *Mem. Inst. Butantan*, *VI Reunião Científica Anual* **61**, 66.
- Hoge, A.R. (1952). Notas herpetológicas. *Mem. Inst. Butantan* **24**(2), 237–240.
- Hoge, A.R. & Belluomini, H.E. (1957/58). Aberrações cromáticas em serpentes brasileiras. *Mem. Inst. Butantan*, **28**, 95–98.
- Melgarejo, A.R. (2003). Serpentes Peçonhentas do Brasil. In: *Animais Peçonhentos no Brasil – Biologia, Clínica e Terapêutica dos Acidentes*. Eds. Cardoso, J.L.C.; França, F.O.S.; Wen, F.H.; Málaque, C.M.S. & Haddad Jr., V. Cap. 4, pp. 33–61. Ed. Sarvier.
- Sazima, I. & Di-Bernardo, M. (1991). Albinismo em Serpentes Neotropicais. *Mem. Inst. Butantan* **53**, 167–173.
- Silva, R.J.; Fontes, M.R.M.; Rodrigues, R.R., Bruder, E.M.; Stein, M.F.B.; Sipoli, G.P.M.; Pinhão, R. & Lopes, C.A.de M. (1999). *J. Venom. Anim. Toxins*, **15** (1).

SILVIA REGINA TRAVAGLIA CARDOSO (E-mail: silviacardoso@butantan.gov.br) and ANA CAROLINA PARPINELLI (E-mail: acp_venom@butantan.gov.br).


Laboratório de Herpetologia, Instituto Butantan, Av. Vital Brazil, 1500, CEP 05503-900, São Paulo, SP, Brazil.

HELICOPS LEOPARDINUS (Water snake):

REPRODUCTION. *Helicops leopardinus* is an aquatic and viviparous snake (Lira-da-Silva *et al.* 1994), distributed from the Guianas to Argentina (Peters & Orejas-Miranda, 1986). It is widespread in Brazil, and one of the most common species in the Pantanal region (Strüssmann & Sazima, 1993). Data on reproduction of this species are scarce. Lira-da-Silva *et al.* (1994) and Freitas (1999) recorded litters ($n = 4$) of seven to 15 newborns for *H. leopardinus* from Bahia, northeast Brazil. Here we present a new record of a litter, as well as provided additional information on the fecundity of the species.

One female *H. leopardinus* collected in the city of Lageado (09°45'S, 48°12'W), Tocantins, northern Brazil, was brought to Instituto Butantan on 21st January 2001. The female (IB 66413; 620 mm in snout-vent length (SVL), 145 mm in tail length (TL), and a mass of 145 g after parturition) gave birth to 31 newborns (21 alive and 10 dead but fully developed) on 8th February 2001. The live newborns averaged 118 mm in SVL (range = 100–135 mm), 39 mm in TL (range = 35–45 mm), and 1.8 g (range = 1.4–2.4 g). The relative clutch mass (RCM; total clutch mass/body mass of mother after parturition; see Shine, 1980) was 0.39. The RCM calculated following Seigel & Fitch (1984) (total clutch mass/body mass of mother + clutch mass) was 0.28.

Figure 1. Relationship between the litter size (including oviductal embryos) and the female's body size in *Helicops leopardinus*. $r_{\text{Spearman}} = 0.82$, $n = 7$, $p = 0.02$.



IB	Region	Co-ordinates	Female SVL (mm)	Litter size
22667	–	–	553	07
20024	Fortaleza, Ceará	03°43'S, 38°32'W	668	25
2646*	Bahia	–	575	19
24534*	Batovi, Mato Grosso	15°51'S, 53°30'W	615	17
44051*	Campo Grande, Mato Grosso do Sul	20°26'S, 54°38'W	545	16
45883*	Campo Grande, Mato Grosso do Sul	20°26'S, 54°38'W	565	12
66413	Lageado, Tocantins	09°45'S, 48°12'W	620	31

Table 1. Summary of data on litter and female's body size for *Helicops leopardinus*. * = oviductal embryos. All preserved individuals were from the Herpetological Collection of the Instituto Butantan (IB).

Additional litters of six preserved specimens presented seven to 25 newborns or oviductal embryos (Table 1). The average litter size in *H. leopardinus* including oviductal embryos is 18 (range = 7–31), and is positively correlated with the female body size ($r_{\text{Spearman}} = 0.82$, $n = 7$, $p = 0.02$; Figure 1). This correlation is a common trend encountered in various snake species (Shine, 1994). Moreover, *H. leopardinus* produce litters of similar size to other congeneric species, particularly *H. infrataeniatus* (Aguiar & Di-Bernardo, 2005) and *H. modestus* (RRS, unpubl. data). The RCM value reported for *Helicops leopardinus* is high but similar to the maximal values reported for other unrelated viviparous and aquatic snakes (e.g. some natricines of the genus *Nerodia* and *Regina*; see Seigel & Fitch, 1984).

ACKNOWLEDGEMENTS

We thank Valdir J. Germano for technical support and Ricardo J. Sawaya for the manuscript review. Finances were supported by FAPESP.

REFERENCES

- Aguiar, L. F. S. & Di-Bernardo, M. (2005). Reproduction of the water snake *Helicops infrataeniatus* Jan, 1865 (Colubridae) in southern Brazil. *Amphibia-Reptilia* **26**, 527–533.
- Freitas, M. A. (1999). Serpentes da Bahia e do Brasil. Feira de Santana: Ed. Dall.
- Lira-da-Silva, R. M. Casais-e-Silva, L. L. Queiroz, I. B. & Nunes, T. B. (1994). Contribuição à biologia de serpentes da Bahia, Brasil. I. Vivíparas. *Rev. Bras. Zool.* **11**, 187–193.
- Peters, J. A. & Orejas-Miranda, B. (1986). Catalogue of Neotropical Squamata. Washington and London: Smithsonian Institution Press.
- Seigel, R. A. & Fitch, H. S. (1984). Ecological patterns of relative clutch mass in snakes. *Oecologia* **61**, 293–301.
- Shine, R. (1980). “Costs” of reproduction in reptiles. *Oecologia* **46**, 92–100.
- Shine, R. (1994). Allometric patterns in the ecology of Australian snakes. *Copeia* **1994**, 851–867.
- Strüssmann, C. & Sazima, I. (1993). The snake assemblage of the Pantanal at Poconé, western Brazil: faunal composition and ecological summary. *Stud. Neotr. Fauna Envir.* **28**, 157–168.
- RODRIGO ROVERI SCARTOZZONI (E-mail: rodrigobuta@yahoo.com.br) and SELMA MARIA DE ALMEIDA-SANTOS (E-mail: almeidasantos@butantan.gov.br).
- Laboratório de Herpetologia, Instituto Butantan, Av. Vital Brazil, 1500, CEP 05503-900, São Paulo, SP, Brazil.